

Rolipa

ABG		Standort	BL	Jahr	Boden / Klima	abs. Ertrag	Sortenmittel	Pflanzenlänge	Lager	Fallzahl	Braunrost	TKM	hl-Gewicht	Verkleisterungs maximum	Verkleisterungs temp	Rhynchosporiu m	
Nr.	Bez.	Ort				dt/ha	dt/ha	cm	Note 1-9	s	Note 1-9	g	kg	AE	°C	Note 1-9	
6	Ackerbau gebiete Süd / Höhenlagen Süd-West	KI-D	BW	2008		27,9	35,2	142	1,0	319	1,0			1480		3,3	
		KI-O	BW	2009		47,5	48,7	136				39,0	76,9				
		KI-C	BW	2011		24,7	35,8	108	1,0								
		HH	BW	2008		55,9	63,3	170	3,7								7,0
				2009		46,0	45,2										
				2010		46,8	55,5			326				1620	76,0		
				2011		44,6	53,4	171	5,0		6,7						
		MHB	BW	2011		32,2	61,7		1,0								
Roggenversuch K.J.Müller 2008																	
		Tosterglope	NI	2008	54 BP; humos, sL	34,2	36,7	166	1,0	174	1,0	33,7				1,0	
		Dottenfelderhof	HE	2008	49 BP; IS	57,6	55,6	180	2,0 bis 7,0	320	7,0	27,9				4,0	
		Höllwangen	BW	2008	60 BP; sL	24,4	37,4	170	5,0	280	1,0	32,5				2,3	